

# Regulation of Group I Metabotropic Glutamate Receptors by MAPK/ERK in Neurons

Li-Min Mao<sup>1</sup> and John Q. Wang<sup>1,2,\*</sup>

<sup>1</sup>Department of Biomedical Sciences, <sup>2</sup>Department of Anesthesiology, School of Medicine, University of Missouri-Kansas City, Kansas City, MO 64108, USA

**Group I metabotropic glutamate receptors (mGluR1 and mGluR5 subtypes) are regulated by protein kinases. A recent focus is mitogen-activated protein kinases (MAPK). A prototypic subclass of MAPKs, extracellular signal-regulated kinases (ERK), is densely expressed in adult brain postmitotic neurons. This kinase resides in not only the cytoplasm around the nucleus, also the neuronal peripheral structures such as synapses. Recombinant ERK2 binds to C terminal tails of mGluR1a *in vitro* and native ERK1/2 forms complexes with mGluR1/5 in neurons *in vivo*. Association of ERK with mGluR1/5 enables the kinase to phosphorylate mGluR1/5 at a cluster of serine sites in the distal C terminus, including a serine residue within the Homer binding site. The ERK-mediated phosphorylation of mGluR1/5 promotes surface expression of mGluR1a in cerebellar neurons. ERK also regulates mGluR1/5 signaling and functions. Among different functional outputs surveyed, ERK exerts an output-specific role in either potentiating or inhibiting their activities. In sum, synaptic group I mGluRs are sufficient substrates of MAPK/ERK. Phosphorylation of mGluR1/5 by ERK has a significant impact on subcellular expression and function of phospho-modified receptors.**

mGluR | striatum | cerebellum | hippocampus | synapse | phosphorylation | G protein-coupled receptors

## Introduction

Glutamate is a key neurotransmitter in the mammalian brain. This transmitter interacts with both ionotropic and metabotropic glutamate receptors (mGluR) to achieve its action. mGluRs are a family of G protein-coupled receptors (GPCR). Base on post-receptor signaling pathways, eight mGluR subtypes are grouped into three functional groups (I, II, and III) (1). Group I mGluRs (mGluR1 and mGluR5 subtypes) have been most extensively investigated. These mGluR subtypes are coupled to phospholipase C $\beta$ 1 (PLC $\beta$ 1) through G $_{\alpha q}$  proteins. Activation of mGluR1/5 increases PLC $\beta$ 1-mediated phosphoinositide hydrolysis, yielding diacylglycerol and inositol-1,4,5-triphosphate (IP $_3$ ) to trigger Ca<sup>2+</sup> and protein kinase C (PKC) signaling pathways, respectively. Noticeably, mGluR1/5 are mostly postsynaptic (2,3). Due to their wide spread distribution in brain regions, mGluR1/5 are actively involved in the regulation of cellular and synaptic activities (1,4) and are linked to various neuropsychiatric, neurodegenerative, and cognitive disorders (4,5).

As membrane-bound GPCRs, mGluR1/5 (mainly long-form splice variants such as mGluR1a, mGluR5a, and mGluR5b) possess a C terminus (CT) protruding into the cytoplasm. The CT tail is large in terms of the number of amino acids and renders mGluR1/5 accessibility to various binding partners (6-8). In fact, various mGluR1/5-associated proteins through interacting with mGluR1/5 CT regulate expression and function of the receptors (7,8). Among different sets of mGluR1/5-associated proteins, protein kinases represent an important group. Accumulated evidence indicates that protein kinases directly interact with intracellular domains of mGluR1/5. Via phosphorylating specific

amino acid(s), a given protein kinase vigorously modulates mGluR1/5 in its distribution and function (9,10).

Mitogen-activated protein kinases (MAPK) are serine/threonine kinases which are densely expressed in adult brain postmitotic neurons. These kinases are activated via a sequential event involving Ras/Rac GTPases, MAPK kinase kinases (Raf or MEKK), and MAPK kinases (MEK) (11). There are three subclasses of MAPKs: extracellular signal-regulated kinases (ERK), c-Jun N-terminal kinases/stress-activated protein kinases (JNK/SAPK), and p38 MAPKs (12). While these subclasses have similar biochemical properties in terms of binding to a specific domain and phosphorylating a common proline-directed motif (S/TP) (13), they are different in upstream activators, downstream substrates, binding and phosphorylation motifs, and thus physiological roles. The prototypic MAPK subclass is ERK which has been extensively studied in its pivotal roles in the modulation of synaptic transmission and plasticity (14,15). Especially, ERK has been recently found to be an important kinase phosphorylating and regulating mGluR1/5. This review then summarizes this recent progress.

## ERK binds to mGluR1/5

ERK1 is associated with mGluR5 in HEK293T cells (16). Similarly, in mouse forebrain lysates, ERK1 and ERK2 were coimmunoprecipitated with mGluR5 (16). Another recent study investigated ERK interactions with mGluR1 (17). Both inactive and active forms of ERK2 directly bound to mGluR1a CT in *in vitro* binding assays with recombinant proteins. The binding site in mGluR1a CT resides in a membrane-proximal region of CT. Native ERK-mGluR1a interactions were studied in the adult rat cerebellum wherein mGluR1 but not mGluR5 is expressed abundantly (18,19). Since mGluR1a is functional at synaptic sites, a defined pool of synaptosomal proteins was extracted and used in coimmunoprecipitation to explore ERK-mGluR1a interactions in the synaptic location. ERK2 and to a lesser extent ERK1 was found to form complexes with mGluR1a in cerebellar neurons under normal conditions (17). In the complexes, the phosphorylated ERK (pERK, i.e., active form of ERK) was found to be predominantly the pERK2 isoform. Thus, ERK (primarily ERK2) is an mGluR1a-associated protein in cerebellar neurons and their interactions could occur at synaptic sites. Of note, ERK is traditionally viewed as a cytoplasmic protein. Once activated, ERK translocates to the nucleus from the cytoplasm to phosphorylate defined transcription factors in order to transcriptionally regulate synaptic transmission and plasticity (14,15,20). In addition, ERK and pERK are present in neuronal peripheral dendritic spines and synaptic zones (21-24). All MAPK cascade components colocalize with ERK2 within the post-

Conflict of Interest: No conflicts declared.

\* Corresponding Author: John Q. Wang, Ph.D.

E-mail: wangjq@umkc.edu

© 2016 by the Authors | Journal of Nature and Science (JNSCI).

synaptic density (PSD) (25,26). Thus, in addition to the nucleus, the synapse is a site where ERK resides and functions to regulate local substrates.

### ERK phosphorylates mGluR1/5

ERK is a proline-directed kinase, meaning that it phosphorylates a consensus motif of serine-proline (SP) or threonine-proline (TP). mGluR1 and 5 both contain a conserved Homer-binding domain (-PPSPF-) in which SP (S1154 for mGluR1a and S1126 for mGluR5) is a potential phosphorylation site subjected to ERK. In fact, using a phospho- and site-specific anti-mGluR5-S1126 antibody, Hu et al. found that expression of constitutively active MEK, which activates ERK, induced mGluR5 phosphorylation in HEK293T cells (16). In cultured cortical neurons, basal immunoreactivity was detected by the anti-mGluR5-S1126 antibody, which was reduced by the MEK inhibitor U0126. Similarly, in adult rat brain tissue, a phospho-specific antibody demonstrated to recognize both mGluR1a and mGluR5 at the Homer binding site (ALTPPpSPFRD) detected phosphorylation signals (27). Phosphorylation was revealed by this antibody in cerebellar mGluR1a of wild type but not mGluR1a knockout mice. Thus, ERK seems to endogenously and dynamically phosphorylate mGluR1/5 at the Homer binding site under normal conditions.

A more detailed phosphorylation mapping study was reported recently. In this study (17), ERK2 strongly phosphorylated the distal part of the GST-fusion mGluR1a CT fragment *in vitro*, while ERK2 did not phosphorylate GST alone and three intracellular loops of mGluR1a (IL1, IL2, and IL3). The ERK2-induced mGluR1a CT phosphorylation was dephosphorylated by a phosphatase. In a site-directed mutation study, a cluster of serine residues in distal CT region were found to be the preferred phosphorylation sites. Noticeably, some phosphorylation sites align well with the consensus ERK phosphorylation motif (PXpSP). When using a motif-specific anti-PXpSP antibody, phosphorylation was found in recombinant mGluR1a CT and in native mGluR1a immunopurified from the synaptosomal pool of the rat cerebellum. The PXpSP immunoreactivity from cerebellar synaptic mGluR1a was reduced by U0126, indicating that ERK normally phosphorylates mGluR1a. The phosphorylation sites are notably different from the location where ERK2 binds to mGluR1a CT. Consistent with this, ERK binds and phosphorylates separated sites in most if not all ERK substrates (13). Interactions between ERK2 and mGluR1a are considered essential for the following phosphorylation step. Inhibition of ERK by U0126 reduced both ERK2-mGluR1a interactions and ERK-mediated phosphorylation of mGluR1a in rat cerebellar neurons.

Phosphorylation of mGluR5-S1126 could be an activity-dependent event. In mouse cultured cortical neurons, applying an mGluR1/5 agonist DHPG for 30 min increased mGluR5-S1126 phosphorylation (16). Since DHPG activated ERK (28), it is thought that the mGluR1/5 agonist activates ERK to trigger a phosphorylation-dependent homologous feedback pathway. Brain-derived neurotrophic factor (BDNF) is also known to activate ERK (29). Adding BDNF elevated mGluR5-S1126 phosphorylation in mouse cultured cortical neurons (16). BDNF produced the same effect in mouse cultured striatal neurons, which was blocked by U0126 (30). The dopamine receptor indirect agonist cocaine and a dopamine D1 receptor agonist SKF38393 increased S1126 phosphorylation in the adult mouse striatum and in cultured striatal neurons, respectively (30). Thus, BDNF and likely D1 signals trigger an ERK-sensitive heterologous signaling pathway to activity-dependently modulate group I mGluRs.

### Regulation of surface expression of mGluR1/5 by ERK

Several mGluR1/5-associated proteins, such as Tamalin, calcineurin inhibitor protein and Norbin, modify expression of mGluR1/5 on the cell surface membrane (31-33). S1126-phosphorylated mGluR5 was enriched in the cell surface of mouse cultured striatal neurons (30). ERK has been found to play a supporting role in constitutive expression of mGluR1a in surface membranes under normal conditions. This was evidenced by the finding that inhibition of ERK by U0126 reduced the level of surface-expressed mGluR1a in rat cerebellar neurons as detected by a surface receptor crosslinking assays *in vivo* (17). It is unclear whether and how ERK-mediated phosphorylation of mGluR1a contributes to this event. Given the fact that the loss of surface-expressed mGluR1a was not accompanied by a concurrent increase in the intracellular pool of the receptor, the subtraction of surface mGluR1a may result from a decrease in synthesis and/or an increase in degradation of the surface receptor.

Phosphorylation of mGluR5 at the Homer binding site increased recombinant Homer binding to mGluR5 peptides containing the Homer binding site *in vitro* (27) and mediates the Preso1-induced Homer binding to mGluR5 in HEK293T cells (16). However, no change in mGluR5 surface expression was observed in *Homer1<sup>-/-</sup>Homer2<sup>-/-</sup>Homer3<sup>-/-</sup>* genotype (34).

### Regulation of mGluR1/5 signaling by ERK

In mouse cultured dorsal spinal cord neurons, sustained treatment with glutamate combined with an NMDA receptor blocker APV and an AMPA receptor blocker NBQX to selectively stimulate mGluR induced a second and delayed  $Ca^{2+}$  rise in addition to an initial  $Ca^{2+}$  rise (16). This delayed  $Ca^{2+}$  response occurred in a higher percentage of neurons after adding U0126 at 2 min after onset of glutamate stimulation, indicating an ERK- and activity-dependent inhibition of mGluRs at least in this functional output pathway (the delayed  $Ca^{2+}$  rise) in these neurons. It is likely that ERK phosphorylation of  $Ca^{2+}$ -coupled mGluR1 or mGluR5 or both contributes to this negative feedback regulation, although direct evidence is lacking. However, ERK phosphorylation of mGluR1/5 may not inhibit all mGluR1/5-mediated outputs. Thus, the role of ERK in regulating mGluR1/5 signaling could be output specific. Indeed, Homer modulates group I mGluR signaling in an output-specific manner. While Homer binding inhibited the delayed  $Ca^{2+}$  responses of group I mGluRs in mouse cultured dorsal spinal horn neurons (16), Homer crosslinking increased mGluR1 coupling to intracellular  $IP_3$  receptors in adult rat cerebellar neurons (35).

In cerebellar neurons, U0126 reduced the DHPG-stimulated  $IP_3$  formation (17). A Tat-fusion peptide that selectively disrupted ERK2 interactions with mGluR1a also produced the similar effect (17). Thus, ERK2 activity is important for the efficacy of mGluR1 signaling in stimulating  $IP_3$  formation.

### Regulation of other mGluR1/5 functions by ERK

ERK phosphorylation of mGluR1/5 may have impact on other mGluR1/5-mediated functions. Whole-cell voltage-clamp recordings identified a slow inward current (SIC) which was an mGluR5-dependent NMDA current in response to DHPG in mouse cultured striatal neurons (30). The recording of this mGluR5-SIC was then used to test mGluR5 function. It was observed that BDNF and a D1 receptor agonist SKF38393 potentiated the mGluR5-SIC, indicating that the two agents may trigger a heterologous mechanism to potentiate mGluR5 signaling. The effect of BDNF was dependent upon ERK because U0126 blocked the BDNF-induced potentiation of the mGluR5-SIC. Moreover, an immediate-early gene form of Homer,

Homer1a, which is usually induced activity-dependently in response to changing synaptic input, plays an important role. It is likely that Homer1a interrupts the binding between constitutively expressed long forms of Homer (Homer1b/c, Homer2, and Homer3) and mGluR5 and thereby facilitates Pin1 to bind to and catalyze isomerization of ERK-phosphorylated mGluR5, resulting in an enhancement of the mGluR5-SIC (30).

S1126 phosphorylation of mGluR1/5 is essential for cocaine sensitization. Repeated administration of cocaine causes a greater motor response to subsequent cocaine exposure, i.e., behavioral sensitization, an animal model of drug addiction. Acute administration of cocaine induced phosphorylation of mGluR5 at S1126 in the striatum of adult mice *in vivo* (30). Behavioral sensitization in response to repeated cocaine was markedly reduced in mice that express mutant mGluR5 which cannot be phosphorylated at S1126. Additionally, depotentiation of corticostriatal long-term potentiation (LTP) occurred in corticostriatal synapses (36). This depotentiation was absent in slices prepared from rodents treated with repeated cocaine administration. Thus, failure of the depotentiation was proposed to be a synaptic correlate of cocaine-induced behavioral sensitization (37,38). Indeed, the D1 receptor agonist SKF38393 inhibited the depotentiation in wild type slices, while SKF38393 did not in slices derived from mutant mice deficient S1126 phosphorylation (30). Thus, mGluR5 phosphorylation at S1126 is required for dopamine inhibition of the depotentiation of corticostriatal LTP, which may serve as a metaplastic basis for motor sensitization.

## Conclusions

Like ionotropic glutamate receptors, group I mGluRs are subject to the regulation by a phosphorylation-dependent mechanism. A recent focus is the MAPK/ERK which resides in not only the nucleus but also peripheral structures such as dendritic synapses. The synaptic sub-pool of ERK has been found to regulate a set of local substrates, including scaffolding proteins (PSD-95 and PSD-93) (39,40), cadherin-associated proteins ( $\delta$ -catenin and plakophilin) (41), and Kv4.2 potassium channels (42,43). Group I mGluRs are another group of substrates of MAPK/ERK at the receptor level. ERK has been found to directly bind to mGluR1

and ERK-mGluR1 interactions are deemed to occur at synaptic sites. Synapse-enriched ERK2 phosphorylates mGluR1/5 at a Homer binding site as well as several other serine sites in the distal region of mGluR1/5 CT. Such phosphorylation is constitutively active and can be upregulated activity-dependently by changing synaptic input. Through phosphorylation, ERK modulates trafficking and expression of mGluR1/5 and controls mGluR1/5 signaling. This ERK-mGluR1/5 coupling in defined sets of synapses undergoes plastic changes in response to chronic disease models and plays an essential role in enduring synaptic and behavioral plasticity. While the studies on ERK and mGluR1/5 have made progresses, future studies are needed to define the specificity of ERK isoforms in phosphorylating specific sites and regulating a distinctive functional output of mGluR1 and mGluR5. Moreover, the role of the ERK-mGluR pathway in a specific brain circuit in a given neuropsychiatric or neurological disorder needs further investigation in the future. In addition to ERK, other subfamilies of MAPKs (JNK and p38) have potential to phosphorylate and regulate mGluR1/5. It will be intriguing to expand the study to these subfamilies to illustrate their roles in comparison with ERK.

If JNK and/or p38 are also involved in the regulation of mGluR1/5, crosstalk among ERK, JNK and p38 may exist. In addition, MAPKs may work in concert with other common protein kinases, such as protein kinase A, PKC and Ca<sup>2+</sup>/calmodulin-dependent protein kinase II, to modulate mGluR1/5. In fact, these kinases have been demonstrated to regulate mGluR1/5 (10). Besides phosphorylation, other types of posttranslational modifications may participate in the regulation of mGluR1/5. Ubiquitination occurs in mGluR1a/5 (44). Palmitoylation and sumoylation take place in ionotropic glutamate receptors (10). Future studies need to explore other types of posttranslational modifications of mGluR1/5 and to dissect temporospatial and functional crosstalk among different types of modifications of the receptors.

## Acknowledgements

This work was supported by NIH grants DA10355 (J.Q.W.) and MH61469 (J.Q.W.).

- Niswender CM, Conn PJ. Metabotropic glutamate receptors: physiology, pharmacology, and disease. *Annu Rev Pharmacol Toxicol.* 2010;50:295-322.
- Lujan R, Nusser Z, Roberts JD, Shigemoto R, Somogyi P. Perisynaptic location of metabotropic glutamate receptors mGluR1 and mGluR5 on dendrites and dendritic spines in the rat hippocampus. *Eur J Neurosci.* 1996; 8:1488-1500.
- Kuwajima M, Hall RA, Aiba A, Smith Y. Subcellular and subsynaptic localization of group I metabotropic glutamate receptors in the monkey subthalamic nucleus. *J Comp Neurol.* 2004;474:589-602.
- Traynelis SF, Wollmuth LP, McBain CJ, Menniti ES, Vance KM, Ogden KK, Hansen KB, Yuan H, Myers SJ, Dingledine R. Glutamate receptor ion channels: structure, regulation, and function. *Pharmacol Rev.* 2010;62:405-496.
- Nicoletti F, Bockaert J, Collingridge GL, Conn PJ, Ferraguti F, Schoepp DD, Wroblewski JT, Pin JP. Metabotropic glutamate receptors: from the workbench to the bedside. *Neuropharmacology.* 2011;60:1017-1041.
- Enz R. The trick of the tail: protein-protein interactions of metabotropic glutamate receptors. *Bioessays.* 2007;29:60-73.
- Enz R. Metabotropic glutamate receptors and interacting proteins: evolving drug targets. *Curr Drug Targets.* 2012;13:145-156.
- Fagni L. Diversity of metabotropic glutamate receptor-interacting proteins and pathophysiological functions. *Adv Exp Med Biol.* 2012;970:63-79.
- Kim CH, Lee J, Lee JY, Roche KW. Metabotropic glutamate receptors: phosphorylation and receptor signaling. *J Neurosci Res.* 2008;86:1-10.
- Mao LM, Guo ML, Jin DZ, Fibuch EE, Choe ES, Wang JQ. Posttranslational modification biology of glutamate receptors and drug addiction. *Front Neuroanat.* 2011;5:19.
- Volmat V, Pouyssegur J. Spatiotemporal regulation of the p42/p44 MAPK pathway. *Biol Cell.* 2001;93:71-79.
- Gallo KA, Johnson GL. Mixed-lineage kinase control of JNK and p38 MAPK pathways. *Nat Rev Mol Cell Biol.* 2002;3:663-672.
- Songyang Z, Lu KP, Kwon YT, Tsai LH, Filhol O, Cochet C, Brickey DA, Soderling TR, Bartleson C, Graves DJ, DeMaggio AJ, Hoekstra MF, Blenis J, Hunter T, Cantley LC. A structure basis for substrate specificities of protein Ser/Thr kinases: primary sequence preference of casein kinases I and II, NIMA, phosphorylase kinase, calmodulin-dependent kinase II, CDK5, and Erk1. *Mol Cell Biol.* 1996;16:6486-6493.
- Sweatt JD. Mitogen-activated protein kinases in synaptic plasticity and memory. *Curr Opin Neurobiol.* 2004;14:311-317.
- Thomas GM, Haganir RL. MAPK cascade signaling and synaptic plasticity. *Nat Rev Neurosci.* 2004;5:173-183.
- Hu JH, Yang L, Kammermeier PJ, Moore CG, Brakeman PR, Tu J, Yu S, Petralia RS, Li Z, Zhang PW, Park JM, Dong X, Xiao B, Worley PF. Preso1 dynamically regulates group I metabotropic glutamate receptors. *Nat Neurosci.* 2012;15:836-844.

17. Yang JH, Mao LM, Choe ES, Wang JQ. Synaptic ERK2 phosphorylates and regulates metabotropic glutamate receptor 1 *in vitro* and in neurons. *Mol Neurobiol*. 2017; in press.
18. Martin LJ, Blackstone CD, Haganir RL, Price DL. Cellular localization of a metabotropic glutamate receptor in rat brain. *Neuron*. 1992;9:259-270.
19. Shigemoto R, Nomura S, Ohishi H, Sugihara H, Nakanishi S, Mizuno N. Immunohistochemical localization of a metabotropic glutamate receptor, mGluR5, in the rat brain. *Neurosci Lett*. 1993;163:53-57.
20. Wang JQ, Fibuch EE, Mao LM. Regulation of mitogen-activated protein kinases by glutamate receptors. *J Neurochem*. 2007;100:1-11.
21. Ortiz J, Harris HW, Guitart X, Terwilliger RZ, Haycock JW, Nestler EJ. Extracellular signal-regulated protein kinases (ERKs) and ERK kinase (MEK) in brain: regional distribution and regulation by chronic morphine. *J Neurosci*. 1995;15:1285-1297.
22. Boggio EM, Putignano E, Sassoe-Pognetto M, Pizzorusso T, Glustetto M. Visual stimulation activates ERK in synaptic and somatic compartments of rat cortical neurons with parallel kinetics. *PLoS ONE*. 2007;2:e604.
23. Sindreu CB, Scheiner ZS, Storm DR. Ca<sup>2+</sup>-stimulated adenylyl cyclases regulate ERK-dependent activation of MSK1 during fear conditioning. *Neuron*. 2007;53:79-89.
24. Mao LM, Reusch JM, Fibuch EE, Liu Z, Wang JQ. Amphetamine increases phosphorylation of MAPK/ERK at synaptic sites in the rat striatum and medial prefrontal cortex. *Brain Res*. 2013;1494: 101-108.
25. Suzuki T, Okumura-Noji K, Nishida E. ERK2-type mitogen-activated protein kinase (MAPK) and its substrates in postsynaptic density fractions from the rat brain. *Neurosci Res*. 1995;22:277-285.
26. Suzuki T, Mitake S, Murata S. Presence of up-stream and downstream components of a mitogen-activated protein kinase pathway in the PSD of the rat forebrain. *Brain Res*. 1999;840:36-44.
27. Orlando LR, Ayala R, Kett LR, Curley AA, Duffner J, Bragg DC, Tsai LH, Dunah AW, Young AB. Phosphorylation of the homer-binding domain of group I metabotropic glutamate receptors by cyclin-dependent kinase 5. *J Neurochem*. 2009;110:557-569.
28. Gallagher SM, Daly CA, Bear MF, Huber KM. Extracellular signal-regulated protein kinase activation is required for metabotropic glutamate receptor-dependent long-term depression in hippocampal area CA1. *J Neurosci*. 2004;24, 4859-4864.
29. Segal RA, Greenberg ME. Intracellular signaling pathways activated by neurotrophic factors. *Annu Rev Neurosci*. 1996;19, 463-489.
30. Park JM, Hu JH, Milshteyn A, Zhang PW, Moore CG, Park S, Datko MC, Domingo RD, Reyes CM, Wang XJ, Etzkorn FA, Xiao B, Szumlinski KK, Kern D, Linden DJ, Worley PF. A prolyl-isomerase mediates dopamine-dependent plasticity and cocaine motor sensitization. *Cell*. 2013;154:637-650.
31. Kitano J, Kimura K, Yamazaki Y, Soda T, Shigemoto R, Nakajima Y, Nakanishi S. Tamalin, a PDZ domain-containing protein, links a protein complex formation of group I metabotropic glutamate receptors and the guanine nucleotide exchange factor cytohesins. *J Neurosci*. 2002;22:1280-1289.
32. Ferreira LT, Dale LB, Ribeiro FM, Babwah AV, Pampillo M, Ferguson SS. Calcineurin inhibitor protein (CAIN) attenuates Group I metabotropic glutamate receptor endocytosis and signaling. *J Biol Chem*. 2009;284:28986-28994.
33. Wang H, Westin L, Nong Y, Birnbaum S, Bendor J, Brismar H, Nestler E, Aperia A, Flajolet M, Greengard P. Norbin is an endogenous regulator of metabotropic glutamate receptor 5 signaling. *Science*. 2009;326:1554-1557.
34. Hu JH, Park JM, Park S, Xiao B, Dehoff MH, Kim S, Hayashi T, Schwarz MK, Haganir RL, Seeburg PH, Linden DJ, Worley PF. Homeostatic scaling requires group I mGluR activation mediated by Homer1a. *Neuron*. 2010;68:1128-1142.
35. Tu JC, Xiao B, Yuan JP, Lanahan AA, Leoffert K, Li M, Linden DJ, Worley PF. Homer binds a novel proline-rich motif and links group I metabotropic glutamate receptors with IP3 receptors. *Neuron*. 1998;21:717-726.
36. Pasini A, Gubellini P, Bonsi P, Conquet F, Picconi B, Centonze D, Bernardi G, Calabresi P. Metabotropic glutamate receptor 5 mediates the potentiation of N-methyl-D-aspartate responses in medium spiny striatal neurons. *Neuroscience*. 2001;106:579-587.
37. Centonze D, Costa C, Rossi S, Prosperetti C, Pisani A, Usiello A, Bernardi G, Mercuri NB, Calabresi P. Chronic cocaine prevents depotentiation at corticostriatal synapses. *Biol Psychiatry*. 2006;60:436-443.
38. Pascoli V, Turlault M, Luscher C. Reversal of cocaine-evoked synaptic potentiation resets during-induced adaptive behaviour. *Nature*. 2012;481:71-75.
39. Sabio G, Reuver S, Feijoo C, Hasegawa M, Thomas GM, Centeno F, Kuhlendahl F, Leal-Ortiz S, Goedert M, Garner C, Cuenda A. Stress- and mitogen-induced phosphorylation of the synapse-associated protein SAP90/PSD-95 by activation of SAPK3/p38gamma and ERK1/ERK2. *Biochem J*. 2004;380:19-30.
40. Guo ML, Xue B, Jin DZ, Mao LM, Wang JQ. Interactions and phosphorylation of postsynaptic density 93 (PSD-93) by extracellular signal-regulated kinase (ERK). *Brain Res*. 2012;1460:18-25.
41. Edbauer D, Cheng D, Batterton MN, Wang CF, Duong DM, Yaffe MB, Peng J, Shang M. Identification and characterization of neuronal mitogen-activated protein kinase substrates using a specific phosphomotif antibody. *Mol Cell Proteomics*. 2009;8:681-695.
42. Adams JP, Anderson AE, Varga AW, Dineley KT, Cook RG, Pfaffinger PJ, Sweatt JD. The A-type potassium channel Kv4.2 is a substrate for the mitogen-activated protein kinase ERK. *J Neurochem*. 2000;75:2277-2287.
43. Schrader LA, Birnbaum SG, Nadin BM, Ren Y, Bui D, Anderson AE, Sweatt JD. ERK/MAPK regulates the Kv4.2 potassium channel by direct phosphorylation of the pore-forming subunit. *Am J Physiol Cell Physiol*. 2006;290:C852-861.
44. Moriyoshi K, Iijima K, Fujii H, Ito H, Cho Y, Nakanishi S. Seven in absentia homolog 1A mediates ubiquitination and degradation of group I metabotropic glutamate receptors. *Proc Natl Acad Sci USA*. 2004;101:8614-8619.